

UNCLASSIFIED

DTIC FILE COPY

②

SECURITY


AD-A198 004

RT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b. RESTRICTIVE MARKINGS							
2a. SECURITY CLASSIFICATION AUTHORITY NA		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for Public Release; Distribution Unlimited							
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE NA									
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report No. 229		5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR- 88-0786							
6a. NAME OF PERFORMING ORGANIZATION University of North Carolina		7a. NAME OF MONITORING ORGANIZATION AFOSR/NM							
6b. ADDRESS (City, State and ZIP Code) Statistics Department CB #3260, Phillips Hall Chapel Hill, NC 27599-3260		7b. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC 20332-6448							
8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F49620 85 C 0144							
8b. OFFICE SYMBOL (If applicable) NM									
9a. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC		10. SOURCE OF FUNDING NOS. <table border="1"><tr><td>PROGRAM ELEMENT NO. 6.1102F</td><td>PROJECT NO. 2304</td><td>TASK NO. A5</td><td>WORK UNIT NO. Leave blank</td></tr></table>		PROGRAM ELEMENT NO. 6.1102F	PROJECT NO. 2304	TASK NO. A5	WORK UNIT NO. Leave blank		
PROGRAM ELEMENT NO. 6.1102F	PROJECT NO. 2304	TASK NO. A5	WORK UNIT NO. Leave blank						
11. TITLE (Include Security Classification) On functional estimates for ill-posed linear problems									
12. PERSONAL AUTHOR(S) Brigola, R. and Keller, A.									
13a. TYPE OF REPORT Reprint		13b. TIME COVERED FROM 9/87 TO 8/88							
14. DATE OF REPORT (Yr., Mo., Day) April 1988		15. PAGE COUNT 10							
16. SUPPLEMENTARY NOTATION N/A									
17. COSATI CODES <table border="1"><tr><td>FIELD</td><td>GROUP</td><td>SUB. GR.</td></tr><tr><td>XXXXXXXXXXXX</td><td></td><td></td></tr></table>		FIELD	GROUP	SUB. GR.	XXXXXXXXXXXX			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Key words and phrases: N/A	
FIELD	GROUP	SUB. GR.							
XXXXXXXXXXXX									
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Ill-posed linear problems in Hilbert space are considered as stochastic filtering problems. Functional estimates of the signal $x$ are given for the problem $Ax=y=z$ where $A$ is a linear, not necessarily bounded operator between Hilbert spaces and $x, y, z$ are Hilbert space valued random elements. As an application, functional estimates are given explicitly for Radon transformed signals with additive white noise.									
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED							
22a. NAME OF RESPONSIBLE INDIVIDUAL Major Brian W. Woodruff		22b. TELEPHONE NUMBER (Include Area Code) (202) 767-5028							
		22c. OFFICE SYMBOL AFOSR/NM							

# CENTER FOR STOCHASTIC PROCESSES

Department of Statistics  
University of North Carolina  
Chapel Hill, North Carolina



ON FUNCTIONAL ESTIMATES FOR ILL-POSED LINEAR PROBLEMS

R. Brigola

and

A. Keller



Technical Report No. 229

April 1988

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

145. G. Kallianpur and V. Perez-Abreu, Stochastic evolution equations with values on the dual of a countably Hilbert nuclear space, July 86. Appl. Math. Optimization, to appear.
146. B.G. Nguyen, Fourier transform of the percolation free energy, July 86. Probab. Theor. Rel. Fields, to appear.
147. A.M. Hsiao, Distribution of the maximum of a Gaussian process by Monte Carlo, July 86. J. Sound Vibration, 112, 1987, 283-293.
148. T. Norbe G., On the existence of probability measures on continuous semi-lattices, Aug. 86.
149. G. Samorodnitsky, Continuity of Gaussian processes, Aug. 86. Ann. Probability, to appear.
150. T. Hsing, J. Husler, and M.R. Leadbetter, Limits for exceedance point processes, Sept. 86. Prob. Theory and Related Fields, to appear.
151. S. Cambanis, Random filters which preserve the stability of random inputs, Sept. 86. Adv. Appl. Probability, 1987, to appear.
152. O. Kallenberg, On the theory of conditioning in point processes, Sept. 86. Proceedings for the First World Congress of the Bernoulli Society, Tashkent, 1986.
153. G. Samorodnitsky, Local moduli of continuity for some classes of Gaussian processes, Sept. 86.
154. V. Mandrekar, On the validity of Beurling theorems in polydiscs, Sept. 86.
155. R.F. Serfozo, Extreme values of queue lengths in  $M/G/1$  and  $GI/M/1$  systems, Sept. 86.
156. B.G. Nguyen, Gap exponents for percolation processes with triangle condition, Sept. 86. J. Statist. Physics, 49, 1987, 235-243.
157. G. Kallianpur and R. Wolpert, Weak convergence of stochastic neuronal models, Oct. 86. Stochastic Methods in Biology, M. Kimura et al., eds., Lecture Notes in Biomathematics, 70, Springer, 1987, 116-145.
158. G. Kallianpur, Stochastic differential equations in duals of nuclear spaces with some applications, Oct. 86. Inst. of Math. & Its Applications, 1986.
159. G. Kallianpur and R.L. Karandikar, The filtering problem for infinite dimensional stochastic processes, Jan. 87. Proc. Workshop on Stochastic Differential Systems, Stochastic Control Theory & Applications, Springer, to appear.
160. V. Perez-Abreu, Multiple stochastic integrals and nonlinear functionals of a nuclear space valued Wiener process, Oct. 86. Appl. Math. Optimization, 16, 1987, 227-245.
161. R.L. Karandikar, On the Feynman-Kac formula and its applications to filtering theory, Oct. 86. Appl. Math. Optimization, to appear.
162. R.L. Taylor and T.-C. Hu, Strong laws of large numbers for arrays or rowwise independent random elements, Nov. 86. Intern. J. Math. & Math. Sci., to appear.
163. M. O'Sullivan and T.R. Fleming, Statistics for the two-sample survival analysis problem based on product limit estimators of the survival functions, Nov. 86.
164. F. Avram, On bilinear forms in Gaussian random variables, Toeplitz matrices and Parseval's relation, Nov. 86.
165. D.B.H. Cline, Joint stable attraction of two sums of products, Nov. 86. J. Multivariate Anal., to appear.
166. R.J. Wilson, Model fields in crossing theory—a weak convergence perspective, Nov. 86.
167. D.B.H. Cline, Consistency for least squares regression estimators with infinite variance data, Dec. 86.
168. L.L. Campbell, Phase distribution in a digital frequency modulation receiver, Nov. 86.
169. B.G. Nguyen, Typical cluster size for 2-dim percolation processes, Dec. 86. J. Statist. Physics, to appear.
170. H. Oodaira, Freidlin-Wentzell type estimates for a class of self-similar processes represented by multiple Wiener integrals, Dec. 86.
171. J. Nolan, Local properties of index- $\beta$  stable fields, Dec. 86. Ann. Probability, to appear.
172. R. Menich and R.F. Serfozo, Optimality of shortest queue routing for dependent service stations, Dec. 86.
173. F. Avram and M.S. Taqqu, Probability bounds for M-Skorohod oscillations, Dec. 86.
174. F. Moricz and R.L. Taylor, Strong laws of large numbers for arrays of orthogonal random variables, Dec. 86.
175. G. Kallianpur and V. Perez-Abreu, Stochastic evolution driven by nuclear space valued martingales, Apr. 87.
176. E. Merzbach, Point processes in the plane, Feb. 87.
177. Y. Kasahara, M. Maejima and W. Vervaat, Log fractional stable processes, March 87.
178. G. Kallianpur, A.G. Mianee and H. Niemi, On the prediction theory of two parameter stationary random fields, March 87.
179. R. Brigola, Remark on the multiple Wiener integral, Mar. 87.
180. R. Brigola, Stochastic filtering solutions for ill-posed linear problems and their extension to measurable transformations, Mar. 87.
181. G. Samorodnitsky, Maxima of symmetric stable processes, Mar. 87.
182. H.L. Hurd, Representation of harmonizable periodically correlated processes and their covariance, Apr. 87.
183. H.L. Hurd, Nonparametric time series analysis for periodically correlated processes, Apr. 87.
184. T. Mori and H. Oodaira, Freidlin-Wentzell estimates and the law of the iterated logarithm for a class of stochastic processes related to symmetric statistics, May 87.

# On functional estimates for ill-posed linear problems

R. Brigola<sup>\*)</sup> and A. Keller

**Abstract:** Ill-posed linear problems in Hilbert space are considered as stochastic filtering problems. Functional estimates of the signal  $x$  are given for the problem  $Ax + y = z$  where  $A$  is a linear, not necessarily bounded operator between Hilbert spaces and  $x, y, z$  are Hilbert space valued random elements. As an application, functional estimates are given explicitly for Radon transformed signals with additive white noise.

**1. Introduction:** Let  $H_1$  and  $H_2$  be Hilbert spaces and  $A: H_1 \longrightarrow H_2$  a linear operator. By Hadamard's definition, a linear problem  $Ax = z$  is well-posed if a solution exists, is unique and depends continuously on the data  $z \in H_2$ . Otherwise it is called ill-posed.

Deterministic regularization methods for ill-posed problems have been extensively treated in literature, starting from the work of A. N. Tichonov and V. Ya. Arsenin [12]. Further references may be found in [6] or [7], for instance. Since an equation  $Ax = z$  often describes a functional relationship between an unknown state  $x$  and an observation  $z$ , which may be affected with random additive noise, ill-posed problems have also been considered as stochastic filtering problems  $Ax + y = z$ , where  $x, y, z$  are Hilbert space valued random elements. For bounded linear transformations  $A$ , defined everywhere in  $H_1$ , stochastic solutions for these equations, depending on various models for the noise  $y$ , have been studied in the work of J. N. Franklin [2], V. Friedrich and A. Uhlig [3].

In this work, we will consider linear transformations  $A$  which are not necessarily bounded or defined everywhere in  $H_1$ , and obtain estimates for a given class of linear functionals of the signal  $x$ , given an observation  $z = Ax + y$ , where  $y$  is a noise on  $H_2$  with positive definite covariance. Signal and noise will be assumed to be zero-mean, Gaussian, weak random variables on  $H_1$  resp.  $H_2$ .

<sup>\*)</sup> The first author's research has been supported by AFOSR Contract No. ~~F 49620-82-C 0009~~

## 2. $D_1$ -best, linear estimates in an operator class $\Delta$

Let  $H_1$  and  $H_2$  be Hilbert spaces and  $A: D_A \rightarrow H_2$  be a linear operator with domain of definition  $D_A \subset H_1$ , and  $A': D_A \rightarrow H_1$  be its formally adjoint operator, i. e.  $\langle Ax, y \rangle = \langle x, A'y \rangle$  for  $x \in D_A$  and  $y \in D_A$ . (cf. [13]).

Here  $\langle \dots \rangle$  denotes the inner product in  $H_1$  resp.  $H_2$ .

Furthermore, let the signal  $x$  be a zero-mean, Gaussian,  $H_1$ -valued weak random variable (cf. [1]) with covariance operator  $B$ , and the noise  $y$  be a zero-mean, Gaussian,  $H_2$ -valued weak random variable with positive definite covariance operator  $C$ . Signal and noise are assumed to be independent, i. e. to have independent finite dimensional distributions.

Given an observation  $z = Ax - y$ , we look for a functional  $h_2 \in H_2$  of the observation  $z$  to estimate a given functional  $\langle x, g \rangle$ ,  $g \in H_1$ , of the signal. More generally, given a subspace  $D_1$  of  $H_1$ , we look for an estimation operator  $L: H_1 \rightarrow H_2$ , transforming the given functionals in  $D_1$  into functionals in  $H_2$ , such that  $\langle z, Lg \rangle$  is a least squares estimate for  $\langle x, g \rangle$  simultaneously for all  $g \in D_1$ . Therefore we define:

### Definition (2.1):

Let  $\Delta \subset \{L: H_1 \rightarrow H_2 \mid L \text{ linear and } D_1 \subset D_L\}$  be a given class of admissible estimation operators.

1) For  $g \in D_1$ , the risk  $r_g(L)$  of  $L \in \Delta$  is given by

$$r_g(L) := E(|\langle z, Lg \rangle - \langle x, g \rangle|^2).$$

2)  $L_0 \in \Delta$  is called  $D_1$ -best, linear estimator for  $x$  in  $\Delta$  if simultaneously for all  $g \in D_1$  and  $L \in \Delta$  it holds:

$$r_g(L_0) \leq r_g(L).$$

It will become clear from the following, that certain restrictions have to be imposed on the class of admissible estimation operators to obtain well-defined domains of definition for the estimations, since we deal with unbounded operators, generally not defined everywhere in the Hilbert spaces  $H_1$  or  $H_2$ .

Now, let  $\Gamma := BA'(ABA' + C)^{-1}$  on its natural domain of definition. If we denote the covariance operator of  $z$  by  $K := ABA' + C$ , then  $K$  is one-to-one, since  $C$  is positive definite and  $B$  is positive semidefinite. Therefore  $\Gamma$  is well-defined and for its domain of definition  $D_\Gamma$  it holds:

$$D_\Gamma = \text{rg}(K) = K(D_{ABA'}),$$

since  $D_K = D_{ABA'} \subset D_{BA'}$ . Also,  $\Gamma' = K^{-1}AB = (ABA' + C)^{-1}AB$  is well-de-

defined on its natural domain. Here  $rg(K)$  denotes the range of  $K$ .

Lemma (2.2):

Let  $rg(A') \subset D_1 \subset D_{\Gamma}$ . Then the following statements hold:

- 1)  $D_1 \subset D_{AB}$
- 2)  $B(D_1) \subset D_A$
- 3)  $AB(D_1) \subset rg(K)$
- 4)  $D_K = D_A$
- 5)  $\Gamma$  is formally adjoint to  $\Gamma'$  on  $D_1$

Proof:

The statements 1) - 3) follow immediately from the assumption  $D_1 \subset D_{\Gamma}$ . Statement 4) follows from 1) and the assumption  $rg(A') \subset D_1$ , and 5) from the definition of  $\Gamma$  and  $\Gamma'$ .

Theorem (2.3):

Let  $\Gamma' := (ABA' + C)^{-1}AB$  and  $rg(A') \subset D_1 \subset D_{\Gamma}$ .

Then  $\Gamma'$  is the  $D_1$ -best, linear estimator for  $x$  in

$$\Delta := \{ L: D_1 \longrightarrow D_A \mid L \text{ linear and } rg(ABA' + C) \subset D_L \}.$$

Proof:

First, we state that  $\Gamma' \in \Delta$ :

$\Gamma'$  is defined on  $D_1$  and by Lemma (2.2), 3) and 4) we obtain:

$\Gamma'(D_1) = K^{-1}AB(D_1) \subset D_K = D_A$ . The formally adjoint  $\Gamma$  of  $\Gamma'$  is defined on  $rg(K)$ , thus  $\Gamma' \in \Delta$ .

The risk of an arbitrary  $L \in \Delta$  for fixed  $g \in D_1$  is given by

$$\begin{aligned} r_g(L) &= E(|\langle x, g \rangle - \langle z, Lg \rangle|^2) = E(|\langle x, g - A'Lg \rangle|^2) \\ &= E(\langle x, g - A'Lg \rangle \overline{\langle y, Lg \rangle}) = E(\overline{\langle x, g - A'Lg \rangle} \langle y, Lg \rangle) + E(|\langle y, Lg \rangle|^2). \end{aligned}$$

By the assumption that  $x$  and  $y$  are zero-mean and independent, we have

$$r_g(L) = \langle Bg, g \rangle + \langle BA'Lg, A'Lg \rangle - \langle BA'Lg, g \rangle - \langle Bg, A'Lg \rangle + \langle CLg, Lg \rangle.$$

Since by Lemma (2.2),  $rg(A') \subset D_1 \subset D_{AB}$  and  $AB(D_1) \subset rg(K)$ , and since  $rg(K) \subset D_L$ , we obtain:

$$\begin{aligned} r_g(L) &= \langle Bg, g \rangle + \langle ABA'Lg, Lg \rangle + \langle CLg, Lg \rangle - \langle BA'Lg, g \rangle - \langle L'ABg, g \rangle \\ &= \langle Bg, g \rangle + \langle L'KLg, g \rangle - \langle BA'Lg, g \rangle - \langle L'ABg, g \rangle. \end{aligned}$$

Thus, comparing the risk of  $L$  with the risk of  $\Gamma'$ , it follows:

$$\begin{aligned} r_g(L) - r_g(\Gamma') &= \langle L'KLg, g \rangle - \langle \Gamma K \Gamma' g, g \rangle - \langle BA'Lg, g \rangle - \langle L'ABg, g \rangle \\ &= \langle BA' \Gamma' g, g \rangle + \langle \Gamma ABg, g \rangle. \end{aligned}$$

Using  $\Gamma K = BA'$  on  $D_A = D_K$  and  $K \Gamma' = AB$  on  $D_1$ , we have:

$$\begin{aligned} r_g(L) - r_g(\Gamma') &= \langle L'KLg, g \rangle - \langle L'K\Gamma'g, g \rangle - \langle \Gamma KLg, g \rangle + \langle \Gamma K\Gamma'g, g \rangle \\ &= \langle (L' - \Gamma')K(L' - \Gamma')'g, g \rangle ; \end{aligned}$$

eventually we have  $r_g(\Gamma') \leq r_g(L)$ , since  $K$  is positive definite.

Remark (2.4):

The operator class  $\Delta$ , which we have used, is maximal in the following sense:

- 1) For an element  $L \in \Delta$ , the condition  $D_1 \subset D_L$  is necessary to define  $\langle z, Lg \rangle$ , and  $L(D_1) \subset D_A$ , is necessary for  $\langle x, A'Lg \rangle$  to exist for all  $g \in D_1$ .
- 2) For  $L \in \Delta$ , the existence of a formally adjoint  $L'$  with  $D_{L'} \supset \text{rg}(K)$  is used to ensure comparability of the elements of  $\Delta$  with respect to the risks  $r_g$ ,  $g \in D_1$ .

3. Application: Functional estimates for noisy Radon transformed signals

A well-known ill-posed linear problem is Radon's integral equation (cf. [10]). The Radon transform has technically been used in Computational Axial Tomography for the reconstruction of a density function from its integrals along hyperplanes. This application motivates, in spite of known inversion formulas, a stochastic treatment, because one has only finitely many data, which additionally may be affected with measurement errors, and the unknown density is in time randomly dependent on body functions of the patient, for instance slight motions during measuring.

We will use the following notations.

Definition (3.1):

Let  $L^1(\mathbb{R}^n)$  be the space of Lesbegue-integrable functions on  $\mathbb{R}^n$ ,  $n \geq 2$ ,  $f \in L^1(\mathbb{R}^n)$ ,  $S^{n-1}$  the unit sphere in  $\mathbb{R}^n$  and  $H(p, q) := \{ x \in \mathbb{R}^n : \langle x, q \rangle = p \}$ ,  $(p, q) \in \mathbb{R} \times S^{n-1}$ , a hyperplane in  $\mathbb{R}^n$ .

Then the Radon transform  $Rf$  of  $f$  is defined by

$$Rf(p, q) := \int_{H(p, q)} f(x) \, m(dx),$$

where  $dm$  is the  $(n-1)$ -dimensional Lesbegue measure on  $H(p, q)$ .

According to [11], the Radon transform is not a continuous operator on the whole of  $L^2(\mathbb{R}^n)$ , the Hilbert space of square integrable functions.

To use the above concept of functional estimation for the problem  $Rx + y = z$ , we will make the following assumptions:

- i) Let  $H_1 := L^2(\mathbb{R}^n)$  and  $H_2 := L^2(\mathbb{R} \times S^{n-1})$ , endowed with the usual inner

products,  $n \geq 2$  a fixed integer.

- ii) If  $\mathcal{F}(\mathbb{R}^n)$  denotes the Schwartz space of rapidly decreasing functions on  $\mathbb{R}^n$ , we identify  $\mathcal{F}(\mathbb{R}^n)$  with a subspace of  $L^2(\mathbb{R}^n)$ , and consider the Radon transform  $R$  as operator from  $L^2(\mathbb{R}^n)$  into  $L^2(\mathbb{R} \times S^{n-1})$  with  $D_R \supset \mathcal{F}(\mathbb{R}^n)$ , (cf. [8]).
- iii) The observation  $z$  is assumed to be  $z = Rx + y$ , where the signal  $x$  is assumed to be a zero-mean, Gaussian,  $H_1$ -valued random element, which is stationary, i. e. its covariance operator  $B$  is given by:

$$\langle g, Bh \rangle = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} b(\|q - s\|) g(q) \overline{h(s)} dq ds, \quad g, h \in \mathcal{F}(\mathbb{R}^n),$$

for some  $b \in \mathcal{F}(\mathbb{R})$ , (cf. [5]).

- iv) The noise  $y$  is assumed to be zero-mean, Gaussian white noise on  $L^2(\mathbb{R} \times S^{n-1})$ , i. e. a weak random variable with covariance operator  $\sigma^2 I$ , with  $I$  the identity operator on  $L^2(\mathbb{R} \times S^{n-1})$ .
- v) The signal and the noise are assumed to be independent.
- vi) Let the class  $D_1$  of functionals, estimates are asked for, be given by  $\mathcal{F}(\mathbb{R}^n)$ .

We will need the following definitions and relations between Radon, Fourier and Hilbert transforms.

Definition (3.2):

- 1) Let  $c := \frac{1}{2(2\pi)^{n-1}}$
- 2) The multiplication operators  $M_k: \mathcal{F}(\mathbb{R} \times S^{n-1}) \longrightarrow L^2(\mathbb{R} \times S^{n-1})$  resp.  $\bar{M}_k: \mathcal{F}(\mathbb{R}^n) \longrightarrow L^2(\mathbb{R}^n)$  are defined for  $k \geq 1$  by:

$$M_k h(p, q) := |p|^{k-1} h(p, q), \quad (p, q) \in \mathbb{R} \times S^{n-1}, \quad h \in \mathcal{F}(\mathbb{R} \times S^{n-1})$$

$$\bar{M}_k g(x) := \|x\|^{k-1} g(x), \quad x \in \mathbb{R}^n, \quad g \in \mathcal{F}(\mathbb{R}^n)$$

$$3) \quad \Phi(x) := \int_{\mathbb{R}^n} b(\|s\|) e^{-i\langle s, x \rangle} ds, \quad x \in \mathbb{R}^n,$$

i. e.  $\Phi$  is the power spectral density of the signal  $x$  (cf. [5]).

Since  $\Phi \in \mathcal{F}(\mathbb{R}^n)$  and depends only on the norm of its argument,  $\varphi(r) := \Phi(\|x\|)$ ,  $r = \|x\|$ ,  $x \in \mathbb{R}^n$ , is well-defined. Of course,  $\Phi$  and  $\varphi$  are non-negative.



4) The multiplication operator  $M_\Phi: \mathcal{F}(\mathbb{R}^n) \longrightarrow \mathcal{F}(\mathbb{R}^n)$  is defined by:

$$M_\Phi g := \Phi \cdot g, \quad g \in \mathcal{F}(\mathbb{R}^n).$$

5) By  $\hat{H}f(t) := \frac{i}{\pi} \int_{\mathbb{R}} \frac{f(p)}{t-p} dp$ ,  $t \in \mathbb{R}$ ,  $f \in \mathcal{F}(\mathbb{R}^n)$ , the Hilbert transform is de-

noted (cf. [9]).

6) For  $g \in \mathcal{F}(\mathbb{R} \times S^{n-1})$ ,  $(p, q) \in \mathbb{R} \times S^{n-1}$ , the operator  $V$  is defined by:

$$Vg(p, q) := \begin{cases} c \left( \frac{\partial}{i \partial p} \right)^{n-1} g(p, q) & \text{for odd } n \\ c i \hat{H} \left( \frac{\partial}{i \partial p} \right)^{n-1} g(p, q) & \text{for even } n \end{cases}$$

(cf. [8]).

Lemma (3.3):

Let  $R$  be the Radon transform and  $F$  be the Fourier transform on  $L^2(\mathbb{R}^n)$ .

Then the following statements hold:

1) For  $(p, q) \in \mathbb{R} \times S^{n-1}$ ,  $f \in L^1(\mathbb{R}^n) \cap L^2(\mathbb{R}^n)$ ,

$$Rf(p, q) = (2\pi)^{\frac{n}{2}-1} \int_{\mathbb{R}} Ff(r, q) e^{ipr} dr$$

2)  $f(x) = (2\pi)^{-n} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} f(s) e^{i\langle s-x, s' \rangle} ds ds'$ , for  $x \in \mathbb{R}^n$ ,  $f \in L^2(\mathbb{R}^n)$ .

The proof is straight forward and left to the reader.

Lemma (3.4):

1)  $R(\mathcal{F}(\mathbb{R}^n)) \subset \mathcal{F}(\mathbb{R} \times S^{n-1})$

2) The adjoint  $R'$  of the Radon transform  $R$  is defined on  $VR(\mathcal{F}(\mathbb{R}^n))$ , where  $V$  is the differential operator from definition (3.2), and  $R'VR = I$ , the identity on  $\mathcal{F}(\mathbb{R}^n)$ .

For the proof it is referred to [8].

Lemma (3.5):

If  $F_p$  denotes the Fourier transform of a function in the variable  $p \in \mathbb{R}$ , then:

1)  $V = c F_p^{-1} M_n F_p$  on  $\mathcal{F}(\mathbb{R} \times S^{n-1})$

2)  $F_p^{-1} M_n F_p R = R F^{-1} \bar{M}_k F$  on  $\mathcal{F}(\mathbb{R}^n)$

3)  $VR = c R F^{-1} \bar{M}_n F$  on  $\mathcal{F}(\mathbb{R}^n)$

For the proof of assertion 1), it is referred to [8]. Assertion 2) and 3) follow immediately from Lemma (3.3) and assertion 1).

**Lemma (3.6):**

For the covariance operator B of the signal x, it holds:

$$B = F^{-1} M_{\Phi} F \text{ on } \mathcal{F}(\mathbb{R}^n)$$

**Proof:**

For given  $g \in \mathcal{F}(\mathbb{R}^n)$ ,  $h \in L^2(\mathbb{R}^n)$  we have:

$$\begin{aligned} \langle Bg, h \rangle &= \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} b(\|s - q\|) g(q) \overline{h(s)} dq ds \\ &= (2\pi)^{-n} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} b(\|s - q\|) \int_{\mathbb{R}^n} Fg(u) e^{i\langle u, q \rangle} du \int_{\mathbb{R}^n} \overline{Fh(v)} e^{-i\langle v, s \rangle} dv dq ds \\ &= (2\pi)^{-n} \int_{\mathbb{R}^n} \left( \int_{\mathbb{R}^n} \left( \int_{\mathbb{R}^n} Fg(u) \Phi(u) e^{i\langle u - v, s \rangle} du \right) ds \right) \overline{Fh(v)} dv, \end{aligned}$$

where the last equality holds by Fubini's theorem, the well-known substitution rule for integrals, and by definition of  $\Phi$ .

Thus, by Lemma (3.3), 2) we obtain:

$$\langle Bg, h \rangle = \int_{\mathbb{R}^n} Fg(v) \Phi(v) \overline{Fh(v)} dv = \langle M_{\Phi} Fg, Fh \rangle,$$

which proves the assertion.

Now, we can calculate best functional estimates for functionals in  $\mathcal{F}(\mathbb{R}^n)$ , according to theorem (2.3), given the problem  $Rx - y = z$  under the above assumptions:

**Theorem (3.7):**

- 1) The operator  $\Gamma: D_{\Gamma} \longrightarrow L^2(\mathbb{R} \times S^{n-1})$ , defined by  $\Gamma := (RBR' + \sigma^2 I)^{-1} RB$ , gives a  $\mathcal{F}(\mathbb{R}^n)$ -best, linear estimator for x in the sense of definition (2.1) within the class:

$$\Delta := \{ L: \mathcal{F}(\mathbb{R}^n) \longrightarrow \text{rg}(VR) \mid L \text{ linear and } \text{rg}(RBR' + \sigma^2 I) \subset D_L \}.$$

Here, V is the operator from definition (3.2).

- 2) For  $(p, q) \in \mathbb{R} \times S^{n-1}$  and  $g \in \mathcal{F}(\mathbb{R}^n)$ , it holds:

$$\Gamma g(p, q) = (2\pi)^{\frac{n-1}{2}-1} \int_{\mathbb{R}} \frac{|r|^{n-1} \varphi(|r|)}{2(2\pi)^{n-1} \varphi(|r|) + \sigma^2 |r|^{n-1}} Fg(rq) e^{ipr} dr.$$

**Proof:**

Clearly,  $\sigma^2 I$  ( $\sigma > 0$ ) is positive definite, and by Lemma (3.4):

$$\text{rg}(R') = R'VR(\mathcal{F}(\mathbb{R}^n)) = \mathcal{F}(\mathbb{R}^n) = D_1.$$

Thus, we have to show  $\mathcal{D}(\mathbb{R}^n) = D_1 \subset D_\Gamma$ , according to theorem (2.3). Let us state, that:

$$\begin{aligned} RBR' + \sigma^2 I &= RB(VR)^{-1} + \sigma^2 VR(VR)^{-1} = R(B + c\sigma^2 F^{-1} \bar{M}_n F)(VR)^{-1} \\ &= RF^{-1}(M_\Phi + c\sigma^2 \bar{M}_n)F(VR)^{-1} \end{aligned}$$

by Lemma (3.4), (3.5) and (3.6).

Therefore, we have:

$$i) (RBR' + \sigma^2 I) VR(\mathcal{D}(\mathbb{R}^n)) = RF^{-1}(M_\Phi + c\sigma^2 \bar{M}_n)F(\mathcal{D}(\mathbb{R}^n))$$

Since  $\Phi \in \mathcal{D}(\mathbb{R}^n)$ , also the functions  $h_g$  on  $\mathbb{R}^n$ , defined by:

$$h_g(x) := \frac{\Phi(x)}{\Phi(x) + c\sigma^2 \|x\|^{n-1}} g(x), \quad g \in \mathcal{D}(\mathbb{R}^n),$$

are elements of  $\mathcal{D}(\mathbb{R}^n)$ .

Hence, we obtain:

$$ii) M_\Phi F(\mathcal{D}(\mathbb{R}^n)) \subset (M_\Phi + c\sigma^2 \bar{M}_n)F(\mathcal{D}(\mathbb{R}^n)).$$

Therefore, application of  $RF^{-1}$  on both sides, Lemma (3.6) and i) yield:

$$RB(\mathcal{D}(\mathbb{R}^n)) \subset RF^{-1}(M_\Phi + c\sigma^2 \bar{M}_n)F(\mathcal{D}(\mathbb{R}^n)) = (RBR' + \sigma^2 I) VR(\mathcal{D}(\mathbb{R}^n)).$$

Thus, we have shown, that  $RB(\mathcal{D}(\mathbb{R}^n)) \subset \text{rg}(RBR' + \sigma^2 I)$ , which implicates:  $\mathcal{D}(\mathbb{R}^n) \subset D_\Gamma$ .

Since  $D_{R'} = VR(\mathcal{D}(\mathbb{R}^n)) = \text{rg}(VR)$ , the class  $\Delta$  equals that one used in theorem (2.3), from which we now obtain assertion 1).

To prove assertion 2), remark that by the above we have for  $g \in \mathcal{D}(\mathbb{R}^n)$ :

$$\begin{aligned} \Gamma'g &= (RBR' + \sigma^2 I)^{-1} RBg = [RF^{-1}(M_\Phi + c\sigma^2 \bar{M}_n)F(VR)^{-1}]^{-1} RBg \\ &= VRF^{-1}(M_\Phi + c\sigma^2 \bar{M}_n)^{-1} FR^{-1} RBg. \end{aligned}$$

Using Lemma (3.5) and (3.6), we get:

$$\Gamma'g = cRF^{-1} \bar{M}_n (M_\Phi + c\sigma^2 \bar{M}_n)^{-1} M_\Phi Fg.$$

Eventually, the representation of the Radon transform in Lemma (3.3) implicates for  $(p, q) \in \mathbb{R} \times S^{n-1}$ ,  $g \in \mathcal{D}(\mathbb{R}^n)$ :

$$\Gamma g(p, q) = (2\pi)^{\frac{n-1}{2}} \int_{\mathbb{R}} \frac{c \|rq\|^{n-1} \Phi(rq)}{\Phi(rq) + c\sigma^2 \|rq\|^{n-1}} Fg(rq) e^{i r p} dr.$$

$$\Gamma g(p, q) = (2\pi)^{\frac{n-1}{2}} \int \frac{|r|^{n-1} \varphi(|r|)}{2(2\pi)^{n-1} \varphi(|r|) + \sigma^2 |r|^{n-1}} Fg(rq) e^{i r p} dr.$$

Remark (3.8):

- 1) Thus, the theorem says, that the evaluation  $\int \Gamma'g(p, q) \overline{z(p, q)} dp dq$  of the observation  $z = Rx - y$  gives the best estimate for the weighted, total density  $\int g(s) x(s) ds$ : the local density  $x(s_0)$ ,  $s_0 \in \mathbb{R}^n$  can be approximated

using the approximate identity as weight functional  $g \in \mathcal{D}(\mathbb{R}^n)$ . With regard to the representation of  $\Gamma'$  in the proof of theorem (3.7), the estimation error is given by:

$$E(|\langle z, \Gamma'g \rangle - \langle x, g \rangle|^2) = \int_{\mathbb{R}^n} \Phi(x) \left( 1 - \frac{2(2\pi)^{n-1} \Phi(x)}{2(2\pi)^{n-1} \Phi(x) + \sigma^2 \|x\|^{n-1}} \right) \overline{f}g(x) \overline{f}g(x) dx$$

- 2) If the Radon transform is considered as a mapping between weighted  $L^2$ -spaces, deterministic regularizations have been worked out by A. K. Louis [7]. Using the above stochastic filtering method, for this case solutions can also be obtained and be related to the result of Tichonov regularization. This will be done elsewhere.

#### Acknowledgement:

The authors want to express their gratitude to Professor D. Kölzow, University of Erlangen, for the suggestion of the subject and helpful discussions. The first author also thanks the hospitality and support of Professor G. Kallianpur, University of North Carolina at Chapel Hill, where part of the work has been done.

#### References:

- [1] Balakrishnan, A. V.: Applied Functional Analysis; Springer, New York-Heidelberg-Berlin, 1976
- [2] Franklin, J. N.: Well-Posed Stochastic Extensions of Ill-Posed Linear Problems; J. Math. Anal. Appl. 31, 1970, 682-716
- [3] Friedrich, V.; Uhlig, A.: Zur stochastischen Regularisierung linearer Gleichungen in Hilberträumen; Beitr. Num. Math. 7, 1979, 33-48
- [4] Gelfand, I. M.; Graev, M. I.; Vilenkin, N. Ya.: Generalized Functions, Vol. V, Academic Press, New York, 1964
- [5] Gelfand, I. M.; Vilenkin, N. Ya.: Generalized Functions, Vol. IV; Academic Press, New York, 1964
- [6] Groetsch, C. W.: The theory of Tikhonov regularization for Fredholm equations of the first kind; Pitman, Boston, 1984
- [7] Louis, A. K.: Tikhonov-Phillips Regularization of the Radon Transform; Int. Ser. Num. Math., Vol. 73, 211-223

- [8] Ludwig, D.: The Radon Transform on Euclidian Space; Comm. Pure Appl. Math. 19, 1966, 49 - 81
- [9] Natterer, F.: The Mathematics of Computerized Tomography; Teubner, Stuttgart, 1986
- [10] Radon, J.: Über die Bestimmung von Funktionen durch ihre Integralwerte längs gewisser Mannigfaltigkeiten; Ber. Verh. Sächs. Akad. Wiss. Leipzig. Math. Natur. Kl. 69, 1917, 262-277
- [11] Smith, K. T.; Solmon, D. C.: Lower Dimensional Integrability of  $L^2$ -Functions; J. Math. Anal. Appl. 51, 1975, 539-549
- [12] Tichonov, A. N.; Arsenin, V., Ya.: Methods of Solving Ill-Posed Problems; Nauka, Moscow, 1974
- [13] Weidmann, J.: Lineare Operatoren in Hilberträumen; Teubner, Stuttgart, 1976

185. R.F. Serfozo, Point processes, May 87. Operations Research Handbook on Stochastic Processes, to appear.
186. Z.D. Bai, W.Q. Liang and W. Vervaat, Strong representation of weak convergence, June 87.
187. O. Kallenberg, Decoupling identities and predictable transformations in exchangeability, June, 87.
188. O. Kallenberg, An elementary approach to the Daniell-Kolmogorov theorem and some related results, June 87. Math. Nachr., to appear.
189. C. Samorodnitsky, Extrema of skewed stable processes, June 87.
190. D. Muirart, M. Sanz and M. Zakai, On the relations between increasing functions associated with two-parameter continuous martingales, June 87.
191. F. Avram and M. Taqqu, Weak convergence of sums of moving averages in the  $\alpha$ -stable domain of attraction, June 87.
192. M.R. Leadbetter, Harald Cramér (1893-1985), July 87. ISI Review, to appear.
193. R. LePage, Predicting transforms of stable noise, July 87.
194. R. LePage and B.M. Schreiber, Strategies based on maximizing expected log, July 87.
195. J. Rosinski, Series representations of infinitely divisible random vectors and a generalized shot noise in Banach spaces, July 87.
196. J. Szulga, On hypercontractivity of  $\alpha$ -stable random variables,  $0 < \alpha < 2$ , July 87.
197. I. Kuznezova-Sholpo and S.T. Rachev, Explicit solutions of moment problems I, July 87. Probability and Mathematical Statistics, 10, 1989, to appear.
198. T. Hsing, On the extreme order statistics for a stationary sequence, July 87.
199. T. Hsing, Characterization of certain point processes, Aug. 87. Stochastic Proc. Appl. 26, 1987, 297-316.
200. J.P. Nolan, Continuity of symmetric stable processes, Aug. 87.
201. M. Marques and S. Cambanis, Admissible and singular translates of stable processes, Aug. 87.
202. O. Kallenberg, One-dimensional uniqueness and convergence results for exchangeable processes, Aug. 87.
203. R.J. Adler, S. Cambanis and G. Samorodnitsky, On stable Markov processes, Sept. 87.
204. C. Kallianpur and V. Perez-Abreu, Stochastic evolution equations driven by nuclear space valued martingales, Sept. 87.
205. R.L. Smith, Approximations in extreme value theory, Sept. 87.
206. E. Willekens, Estimation of convolution tails, Sept. 87.
207. J. Rosinski, On path properties of certain infinitely divisible processes, Sept. 87.
208. A.H. Koreziloglu, Computation of filters by sampling and quantization, Sept. 87.
209. J. Bather, Stopping rules and observed significance levels, Sept. 87.
210. S.T. Rachev and J.E. Yukich, Convolution metrics and rates of convergence in the central limit theorem, Sept. 87.
211. M. Fujisaki, Normed Bellman equation with degenerate diffusion coefficients and its applications to differential equations, Oct. 87.
212. G. Simons, Y.C. Yao and X. Wu, Sequential tests for the drift of a Wiener process with a smooth prior, and the heat equation, Oct. 87.
213. R.L. Smith, Extreme value theory for dependent sequences via the Stein-Chen method of Poisson approximation, Oct. 87.
214. C. Houdré, A note on vector bimeasures, Nov. 87.
215. M.R. Leadbetter, On the exceedance random measures for stationary processes, Nov. 87.
216. M. Marques, A study on Lebesgue decomposition of measures induced by stable processes, Nov. 87.
217. M.T. Alpuim, High level exceedances in stationary sequences with extremal index, Dec. 87.
218. R.F. Serfozo, Poisson functionals of Markov processes and queueing networks, Dec. 87.
219. J. Bather, Stopping rules and ordered families of distributions, Dec. 87.
220. S. Cambanis and M. Maejima, Two classes of self-similar stable processes with stationary increments, Jan. 88.
221. H.P. Hücke, G. Kallianpur and R.L. Karandikar, Smoothness properties of the conditional expectation in finitely additive white noise filtering, Jan. 88.
222. I. Mitoma, Weak solution of the Langevin equation on a generalized functional space, Feb. 88.
223. L. de Haan, S.I. Resnick, H. Rootzén and C. de Vries, Extremal behaviour of solutions to a stochastic difference equation with applications to arch-processes, Feb. 88.
224. O. Kallenberg and J. Szulga, Multiple integration with respect to Poisson and Lévy processes, Feb. 88.
225. D.A. Dawson and L.C. Corostiza, Generalized solutions of a class of nuclear space valued stochastic evolution equations, Feb. 88.
226. G. Samorodnitsky and J. Szulga, An asymptotic evaluation of the tail of a multiple symmetric  $\alpha$ -stable integral, Feb. 88.
227. J.J. Hunter, The computation of stationary distributions of Markov chains through perturbations, Mar. 88.
228. H.C. Ho and T.C. Sun, Limiting distribution of nonlinear vector functions of stationary Gaussian processes, Mar. 88.
229. R. Brigola, On functional estimates for ill-posed linear problems, Apr. 88.